

SUBJECTS

- Achrysonocharoides* sp.
 Coffee leaf miner parasitoid in Puerto Rico, 72(2), 255.
- Alfalfa, 9 cultivars
 Forage production in semiarid region, 72(2), 221.
- Allelopathic effects of weeds
 Pumpkin, 72(3), 491.
- Amaranthus dubius*
 Allelopathic effect on pumpkin, 72(3), 491.
 Exudates from its roots in eggplants, 72(3), 495.
- Anastrepha obliqua*
 Bacteria associated, 72(3), 489.
- Anastrepha* sp.
 Identity and economic importance, 72(2), 325.
- Anthonomus eugenii*
 Sampling methods for population estimation, 72(3), 387.
 Yield-loss components and economic thresholds, 72(3), 375.
- Anthracnose
 Damage to immature soybean seeds, 72(1), 73.
 Yield and profit in monocultures and interplantings of yams, 72(2), 179.
- Asparagus
 Seed production, 72(4), 619.
- Bacillus coagulans*
 Recovery from food refrigerated before incubation, 72(4), 617.
- Bacillus maurus*
 Recovery from food refrigerated before incubation, 72(4), 617.
- Bacteria
 Recovery from food refrigerated before incubation, 72(4), 617.
- Banana
 Dry matter composition, 72(3), 337.
 Nutrient uptake, 72(3), 337.
- Bean
 Yield response to tillage techniques, 72(1), 119.
 Bean, winged
 Plant spacing and growth and yield, 72(2), 273.
- Beans
 Inoculation with *Rhizobium*, 72(3), 427.
- Beans, dry
 Varying cultural practices at two locations, 72(4), 515.
- Beans, seed yields
 Heritabilities and phenotypic correlations, 72(2), 301.
- Beskia aelopa*
 Natural enemy of pentatomids, 72(3), 371.
- Brachiaria brizantha*
 Production in the humid northern coastal plains, 72(1), 91.
- Brachiaria humicola*
 Production in the humid mountains, 72(1), 99.
- Brachiaria ruiziiensis*
 Production in the humid northern coastal plains, 72(1), 91.
- Café
 Susceptibilidad de cultivares al minador, 72(1), 165.
- Cajanus cajan*
 Determinate genotypes, 72(4), 545.
 Light trap studies of *Heliothis virescens* and *Etiella zinckenella*, 72(4), 557.
 Some host plants of its leafhopper, 72(3), 499.
- Callie*, hay
 Storage and digestibility, voluntary intake and chemical components, 72(4), 531.
- Capsicum annuum*
 Yield on northwestern coast, 72(3), 395.
- Cercospora kikuchii*
 Damage to immature soybean seeds, 72(1), 73.
- Chrysotomota* sp.
 Coffee leaf miner parasitoid in Puerto Rico, 72(2), 255.
- Cirrospilooides* sp.
 Coffee leaf miner parasitoid in Puerto Rico, 72(2), 255.
- Citron, bars
 Freezing and shelf life, 72(1), 171.
- Citrus medica*, bars
 Freezing and shelf life, 72(1), 171.
- Cocoyam, in vitro cultured tissue
 Salt and sugar levels for effective plant regeneration, 72(4), 515.

- Coffee
 Bacterial leaf spot, 72(4), 621.
 Susceptibility of cultivars to leaf miner, 72(1), 165.
- Coffea arabica*
 Distribution of leaf miner and its parasitoids in tree canopy, 72(1), 141.
- Coffee leaf miner
 Distribution in the tree canopy, 72(1), 141.
- Parasitoids in Puerto Rico, 72(2), 255.
- Coleoptera
 A new rice pest for Puerto Rico, 72(2), 333.
- Colletotrichum dematium*
 Damage to immature soybean seeds, 72(1), 73.
- Colletotrichum gloeosporioides*
 Yield and profit in monocultures and interplantings of yams, 72(2), 179.
- Coniothyrium* leaf spot
 A newly described disease of sorghum, 72(2), 319.
- Consumptive use, seasonal
 Bell and Cubanelle peppers, 72(4), 609.
- Contarinia sorghicola*
 A newly introduced pest to Puerto Rico, 72(1), 161.
- Corn
 Yield response to tillage techniques, 72(1), 119.
- Corn, Opaque-2 populations
 Agronomic performance, 72(2), 309.
- Corn, sweet
 Estimation of irrigation requirements, 72(2), 277.
- Corn, two locations
 N applications and population densities, 72(1), 127.
- Cosmopolites sordidus*
 Resistance of Lacknau plantain cultivar, 72(3), 353.
- Cows, dairy, confinement feeding
 Green chopped stargrass or hay, 72(2), 231.
- Crotalaria*
 Susceptibility to *Etiella zinckenella*, 72(1), 147.
- Crotalaria palida*
 Hymenopterous parasitoid of larvae feeding on, 72(1), 153.
- Cucurbita moschata*
 Allelopathic effects of weeds, 72(3), 491.
 Diquat as a postemergence herbicide, 72(2), 285.
- Curculionidae
 A new rice pest for Puerto Rico, 72(2), 333.
- Cynodon dactylon*
 Production in humid northern coastal plains (Barceloneta), 72(2), 201.
 Production in semiarid region, 72(2), 211.
 Production in the humid mountains, 72(1), 99.
 Production in the humid northern coastal plains (Toa Baja), 72(1), 91.
 Response to potassium fertilization, 72(1), 81.
- Cynodon dactylon*, hay
 Storage and digestibility, voluntary intake and chemical components, 72(4), 531.
 Yield and nutritive value at three harvesting intervals, 72(1), 109.
- Cynodon nemfuensis*
 Confinement feeding of dairy cows, 72(2), 231.
 Production in humid northern coastal plains (Barceloneta), 72(2), 201.
 Production in semiarid region, 72(2), 211.
 Production in the humid mountains, 72(1), 99.
 Production in the humid northern coastal plains (Toa Baja), 72(1), 91.
 Response to potassium fertilization, 72(1), 81.
- Cynodon nemfuensis*, hay
 Yield and nutritive value at three harvesting intervals, 72(1), 109.
- Cynodon pectostachyus*
 Production in humid northern coastal plains (Barceloneta), 72(2), 201.
 Production in semiarid region, 72(2), 211.
 Production in the humid mountains, 72(1), 99.
 Production in the humid northern coastal plains (Toa Baja), 72(1), 91.
 Storage and digestibility, voluntary intake and chemical components, 72(4), 531.
- Cynodon pectostachyus*, hay from
 Yield and nutritive value at three harvesting intervals, 72(1), 109.
- Cynodon* sp.
 Production in humid northern coastal plains, 72(2), 201.
- Dichanthium annulatum*
 Production in semiarid region, 72(2), 211.
- Digitaria decumbens*
 Production in the humid mountains, 72(1), 99.
 Production in the humid northern coastal plains (Barceloneta), 72(2), 201.
 Production in the humid northern coastal plains (Toa Baja), 72(1), 91.

- Response to potassium fertilization, 72(1), 81.
- Digitaria milanjiana*
Production in humid northern coastal plains, 72(2), 201.
- Digitaria pentzii*
Production in humid northern coastal plains, 72(2), 201.
Production in the humid mountains, 72(1), 99.
Response to potassium fertilization, 72(1), 81.
- Digitaria pentzii*, hay
Storage and digestibility, voluntary intake and chemical components, 72(4), 531.
Yield and nutritive value at three harvesting intervals, 72(1), 109.
- Dioscorea* spp.
Anthracnose, yield and profit, 72(2), 179.
- Diquat*
Post emergence herbicide in pumpkin, 72(2), 285.
- Dwarfing interstems in trees
Growth and yield components in mango, 72(4), 501.
- Echinochloa colonum*
Allelopathic effect on pumpkin, 72(3), 491.
Exudates from its roots in eggplants, 72(3), 495.
- Ectrichodiinae*
Notes about *Rhiginia* sp., 72(2), 251.
- Eggplant
Yield response to tillage techniques, 72(1), 119.
- Eggplant transplants
Exudates of roots of weeds, 72(3), 495.
- Elevation
Relation to temperature, 72(3), 449.
- Empoasca* spp.
Some host plants, 72(3), 499.
- Endosperm, hard
Performance of Opaque-2 populations in Puerto Rico, 72(2), 309.
- Eriochloa polystachya*
Production in the humid northern coastal plains, 72(1), 91.
- Ethanol
Production from blackstrap molasses by *Zymomonas mobilis* and *Saccharomyces*, 72(3), 488.
- Etiella zinckenella*
Damage to immature soybean seeds, 72(1), 73.
Natural enemies, 72(1), 153.
Susceptibility of *Crotalaria* spp. to its attack, 72(1), 147.
- Etiella zinckenella* on pigeon pea
Light trap studies, 72(4), 557.
- Euphorbia heterophylla*
Allelopathic effect on pumpkin, 72(3), 491.
Exudates from its roots in eggplants, 72(3), 495.
- Evapotranspiration, potential
Hargreaves-Samani technique, 72(1), 57.
Vieques Island, 72(1), 177.
- Fascioliasis
Relationship to watercress cultivation, 72(3), 405.
- Fermentation batch
Patterns for strains of *Zymomonas*, 72(3), 475.
- Fermentation, batch and fed-batch
High test molasses by *Zymomonas mobilis*, 72(3), 485.
- Fermentation, high test molasses
Commercial yeast nutrient, 72(1), 173.
Commercial yeast nutrient evaluation, 72(4), 509.
Compared with blackstrap molasses, 72(1), 9.
- Flies, fruit
Identity and economic importance, 72(2), 325.
- Fly, fruit, West Indian
Bacteria associated, 72(3), 489.
- Fodder, hay, stargrass
Confinement feeding of dairy cows, 72(2), 231.
- Forage, alfalfa
Production of 9 cultivars in semiarid region, 72(2), 221.
- Fruit fly, West Indian
Bacteria associated, 72(3), 489.
- Fungi, vesicular-arbuscular
Inoculation of sour orange seedlings, 72(2), 191.
- Glomus* spp.
Inoculation of sour orange seedlings, 72(2), 191.
- Glyphosate
Optimum time interval and frequency of application, 72(2), 297.
- Grapes, table
Variety evaluation at Fortuna, 72(2), 335.
- Grasses, tropical
Response to potassium fertilization, 72(1), 81.
- Grasses, tropical, grazed
Production in humid mountains, 72(1), 99.
Production in humid northern coastal plains (Barceloneta), 72(2), 201.
Production in humid northern coastal plains (Toa Baja), 72(1), 91.

- Production in semiarid region, 72(2), 211.
- Grasses, tropical, hay
Storage and digestibility, voluntary intake and chemical components, 72(4), 531.
- Grasses, tropical, hay from
Yield and nutritive value at three harvesting intervals, 72(1), 109.
- Guinea grass, hay
Storage and digestibility, voluntary intake and chemical components, 72(4), 531.
- Gymnocalystia*
Natural properties of pentatomids, 72(3), 371.
- Hargreaves and Samani
Model for potential evapotranspiration estimation, 72(1), 177.
- Hargreaves-Samani technique
Potential evapotranspiration for the south coast, 72(1), 57.
- Hay, fodder, stargrass
Confinement feeding of dairy cows, 72(2), 231.
- Hay, tropical grasses
Storage and digestibility, voluntary intake and chemical components, 72(4), 531.
- Yield and nutritive value at three harvesting intervals, 72(1), 109.
- Heliothis virescens* on pigeonpea
Light trap studies, 72(4), 557.
- Hemiptera
Description of new species of *Rasahus*, 72(2), 247.
- Description of *Rhiginia dureti* Carpintero from Peru, 72(2), 251.
- Heritabilities and phenotypic correlations
Bean seed yield and yield components, 72(2), 301.
- Heterospilus etiellae*
Larval parasitoid, 72(1), 153.
- Horismenus* sp.
Coffee leaf miner parasitoid in Puerto Rico, 72(2), 255.
- Iconella etiellae*, larval parasitoid
Recovered from soybean, 72(1), 153.
- Inoculation, sour orange seedlings
Vesicular-arbuscular mycorrhizal fungi, 72(2), 191.
- Inoculum, distillers' dry yeast
Rum fermentation on pilot plant scale, 72(3), 445.
- Interstems in mango
Dwarfing effect on growth and yield, 72(4), 501.
- Ipomoea batatas*
Planting dates, morphology and growth, 72(3), 437.
- Sensory evaluation, 72(2), 265.
- Ipomoea batatas*, bars
Freezing and shelf life, 72(1), 171.
- Irrigation, drip
Cost for vegetable farming, 72(1), 31.
- Labor input in vegetable production, 72(1), 41.
- Root distribution of fertigated peppers, 72(1), 51.
- Irrigation (drip) and plastic mulch
Nutrient uptake in sweet peppers, 72(4), 575.
- Irrigation of sweet corn
Estimation requirements, 72(2), 277.
- Leaf miner, coffee
Distribution of parasitoids in the tree canopy, 72(1), 141.
- Susceptibility of coffee cultivars, 72(1), 165.
- Leaf spot
A *Coniothyrium* disease in sorghum, 72(2), 319.
- Leaf spot, bacteria
Coffee, 72(4), 621.
- Leafhopper, pigeon pea
Some host plants, 72(3), 499.
- Lepidoptera, larvae
Chemical control in tomato, 72(1), 169.
- Leucoptera coffeella*
Distribution of parasitoids in the tree canopy, 72(1), 141.
- Parasitoids in Puerto Rico, 72(2), 255.
- Susceptibility of coffee cultivars, 72(1), 165.
- Lima bean pod borer
Susceptibility of *Crotalaria* spp to its attack, 72(1), 147.
- Liriomyza sativae*
Chemical control in tomato, 72(1), 169.
- Lissorhoptrus isthmicus*
A new rice pest for Puerto Rico, 72(2), 333.
- Lycopersicon esculentum*
Chemical control of *Liriomyza sativae* and Lepidoptera larvae, 72(1), 169.
- Mabí, bark
Chemical constituents, 72(4), 625.
- Maíz dulce
Irrigation requirements, estimation of, 72(2), 277.
- Requisitos de riego, estimación de, 72(2), 277.
- Tres niveles de riego, crecimiento y rendimiento, 72(4), 565.

- Mango
Fruit flies which attack, 72(2), 325.
- Mango, dwarfing rootstocks
Tree size and yield of varieties, 72(1), 1.
- Mango, growth and yield
Dwarfing effect of interstems, 72(4), 501.
- Mangoes, hot water treated
Shelf life and acceptability, 72(3), 469.
- Mangoes, 3 cultivars
Shelf life when hot water-treated, 72(3), 469.
- Medicago sativa*, 9 cultivars
Forage production in semiarid region, 72(2), 221.
- Mimosa*, climbing
Chemical control in a local pasture, 72(4), 615.
- Miraz insularis*
Coffee leaf miner parasitoid in Puerto Rico, 72(2), 255.
Parasitoid of coffee leaf miner, 72(1), 141.
- Molasses, blackstrap
Ethanol production by *Zymomonas mobilis* and *Saccharomyces* sp., 72(3), 488.
- Molasses high test
Batch and fed-batch fermentation by *Zymomonas mobilis*, 72(3), 485.
Commercial yeast nutrient for fermentation, 72(1), 173.
Pilot plant scale evaluation of a commercial yeast nutrient in fermentation, 72(4), 509.
- Molasses, high test, fermentation
Compared with blackstrap, 72(1), 9.
- Molasses, high test vs blackstrap
Batch fermentation patterns for strain of *Zymomonas*, 72(3), 475.
- Mormidea angustata* Stål
Host plants in rice fields, 72(3), 365.
Pentatomid parasitized by natural enemies, 72(3), 371.
- Musa* sp.
Optimum time interval and frequency of glyphosate application, 72(2), 297.
- Mushroom, straw
Carbon and nitrogen nutritional influences on growth, 72(1), 19.
- Nasturtium officinale*
Relationship of cultivation sites to fascioliasis incidence, 72(3), 405.
- Nezara viridula*
Damage to immature soybean seeds, 72(1), 73.
- Nitrogen applications
Corn at two locations, 72(1), 127.
- Oebalus ypsilon-griseus*
Host plants in rice fields, 72(3), 365.
Pentatomid parasitized by natural enemies, 72(3), 371.
- Oencyrtus submetallicus*
Natural enemy of pentatomids, 72(3), 371.
- Opaque-2 corn populations
Agronomic performance, 72(2), 309.
- Orange, sour, seedlings
Inoculation with vesicular-arbuscular mycorrhizal fungi, 72(2), 191.
- Panicum maximum*
Production in humid northern coastal plains (Barceloneta), 72(2), 201.
Production in semiarid region, 72(2), 211.
Production in the humid mountains, 72(1), 99.
Production in the humid northern coastal plains (Toa Baja), 72(1), 91.
Storage and digestibility, voluntary intake and chemical components, 72(4), 531.
- Panicum maximum*, hay
Yield and nutritive value at three harvesting intervals, 72(1), 109.
- Paraquat
Postemergence weed control in pumpkin, 72(2), 285.
- Partenium hysterophorus*
Allelopathic effect on pumpkin, 72(3), 491.
Exudates from its roots in eggplants, 72(3), 495.
- Pentatomids
Host plants in rice fields, 72(3), 365.
Natural enemies in rice fields, 72(3), 371.
- Pepper
Yield response to tillage technique, 72(1), 119.
- Pepper cultivars
Yield on northwestern coast, 72(3), 395.
- Pepper weevil
Sampling methods for population estimation, 72(3), 387.
Yield-loss components and economic thresholds, 72(3), 375.
- Peppers
Seasonal consumptive use of water, 72(4), 609.
- Peppers, sweet
Root distribution of nitrogen fertigated, 72(1), 51.
- Peppers, sweet, nitrogen fertigated
Nutrient uptake and growth characteristics, 72(4), 575.

- Peronospora manshurica*
Damage to immature soybean seeds, 72(1), 73.
- Phaseolus vulgaris*
Heritabilities and phenotypic correlations, 72(2), 301.
Inoculation with *Rhizobium*, 72(3), 427.
Varying cultural practices at two locations, 72(4), 521.
- Phenotypic correlations
Bean seed yield and yield components, 72(2), 301.
- Phomopsis* spp.
Damage to immature soybean seeds, 72(1), 73.
- Pigeon pea
Determinate genotypes, 72(4), 545.
Light trap studies of *Heliothis virescens* and *Etiella zinckenella*, 72(4), 557.
Some host plants of its leafhopper, 72(3), 499.
- Plantain
Irrigation requirements, 72(4), 599.
Optimum time interval and frequency of glyphosate application, 72(2), 297.
- Plantain, Lacknau cultivar
Resistance to corm weevil, 72(3), 353.
- Plátano
Requisitos de riego, 72(4), 599.
- Potassium fertilization
Response of tropical grasses in the humid coastal region, 72(1), 81.
- Pseudomonas syringae*
Leaf spot in coffee, 72(4), 621.
- Psophocarpus tetragonolobus*
Plant spacing, growth, yield, 72(2), 273.
- Puccinia purpurea*
Sorghum, 72(1), 65.
- Pumpkin
Allelopathic effects of weeds, 72(3), 491.
Diquat as a postemergence herbicide, 72(2), 285.
- Rasahus* sp.
Description of new species, 72(2), 247.
- Reduviidae
Description of new species of *Rasahus*, 72(2), 247.
- Description of *Rhiginia dureti* Carpintero from Peru, 72(2), 251.
- Rhiginia* sp.
New species described from Peru, 72(2), 251.
- Rhizobium*
Inoculation of *Phaseolus vulgaris*, 72(3), 427.
- Rice
Attacked by *Lissorhoptrus isthmicus* in Puerto Rico, 72(2), 333.
- New disease record from Puerto Rico, 72(2), 331 (RN).
- Rice fields
Host plants of pentatomids, 72(3), 365.
Natural enemies of pentatomids, 72(3), 371.
- Riego de maíz dulce
Estimación de los requisitos, 72(2), 277.
- Root distribution, sweet peppers
Nitrogen fertigated and drip irrigated, 72(1), 51.
- Rottboelia exaltata*
Allelopathic effect on pumpkin, 72(3), 491.
Exudates from its roots in eggplants, 72(3), 495.
- Rootstocks, dwarfing
Tree size and yield of varieties, 72(1), 1.
- Rum fermentation
Distillers' dry yeast inoculum, 72(3), 445.
- Rust, sorghum, 72(1), 65.
- Saccharomyces*
Ethanol production from blackstrap molasses, 72(3), 483.
- Santainezia bicolor*
Synonym of *Rhiginia haenschi*, 72(2), 251.
- Seed asparagus
Production in Puerto Rico, 72(4), 619.
- Seed, yield and yield components
Heritabilities and phenotypic correlations in beans, 72(2), 301.
- Seedlings, orange, sour
Inoculation with vesicular-arbuscular mycorrhizal fungi, 72(2), 191.
- Shelf-life and acceptability
Hot water-treated mangoes, 72(3), 469.
- Smut, false
A new disease record from Puerto Rico's rice fields, 72(2), 331.
- Solanum melongena*
Exudates of roots of weeds, 72(3), 495.
- Sorghum
A newly described disease caused by *Coniothyrium*, 72(2), 319.
Irrigation requirements in the south and north coasts, 72(4), 585.
- Sorghum halepense*
Allelopathic effect on pumpkin, 72(3), 491.
Exudates from its roots in eggplants, 72(3), 495.
- Sorghum midge
A newly introduced pest to Puerto Rico, 72(1), 161.
- Sorghum rust, 72(1), 65.

- Sorgo**
Requisitos de riego en las costas sur y norte, 72(4), 585.
- Soybean seeds, immature**
Fungi and insect damage, 72(1), 73.
- Spacing**
Growth and yield of winged bean, 72(2), 273.
- Stargrass, green chopped fodder or hay**
Confinement feeding of dairy cow, 72(2), 231.
- Sweet potato, bars**
Freezing and shelf life, 72(1), 171.
- Sweet potato, 4 clones**
Planting dates, morphology and growth, 72(3), 437.
- Sweet potato selections, cooked**
Sensory evaluation, 72(2), 265.
- Tanier**
In vitro propagation, 72(3), 413.
- Temperature**
Relation to elevation, 72(3), 449.
- Thrips palmi**
A new insect pest in Puerto Rico, 72(2), 327.
- Tillage techniques**
Yield response of vegetables in semiarid region, 72(1), 119.
- Tissue culture, cocoyam**
Salt and sugar levels for effective plant regeneration, 72(4), 515.
- Tomate**
Periodo critico de competencia de malezas, 72(2), 291.
- Tomato**
Chemical control of *Liriomyza sativae* and Lepidoptera larvae, 72(1), 169.
Critical period of weed competition, 72(2), 291.
Yield response to tillage technique, 72(1), 119.
- Trianthema portulacastrum**
Allelopathic effect on pumpkin, 72(3), 491.
Exudates from its roots in eggplants, 72(3), 495.
- Trichogramma sp**
Parasitism on *Etiella zinckenella* eggs, 72(1), 153.
- Ustilaginoidea oryzae**
A new disease record from Puerto Rico's rice fields, 72(2), 331.
- Uva de mesa**
Evaluación de variedades, 72(2), 335.
- Vegetable farming**
Cost of drip irrigation system, 72(1), 31.
- Vegetables**
Yield response to tillage technique, 72(1), 119.
- Vegetables, drip irrigated**
Labor input requirements, 72(1), 41.
- Vieques Island**
Potential evapotranspiration with Har-greaves and Samani model, 72(1), 177.
- Vitis vinifera**
Evaluación de variedades, 72(2), 335.
Variety evaluation at Fortuna, 72(2), 335.
- Volvariella volvacea**
Carbon and nitrogen nutritional influences on growth, 72(1), 19.
- Watercress**
Relationship of cultivation sites to fascioliasis incidence, 72(3), 405.
- Watermelon**
Yield response to tillage technique, 72(1), 119.
- Weeds**
Critical competition period in tomatoes, 72(2), 291.
Exudates from weed roots and eggplant transplants, 72(3), 495.
- Weeds, allelopathic effects**
Pumpkin, 72(3), 491.
- Weevil, corm, plantain**
Resistance of Lacknau plantain cultivar, 72(3), 353.
- Weevil, pepper**
Sampling methods for population estimation, 72(3), 387.
Yield-loss components and economic thresholds, 72(3), 375.
- Xanthosoma spp.**
In vitro propagation, 72(3), 413.
- Yam, monocultures and interplantings Anthracnose, yield and profit, 72(2), 179.**
- Yeast, commercial**
Nutrient for high test molasses fermentation, 72(1), 173.
- Yeast, distillers' dry**
Inoculum for rum fermentation, 72(3), 445.
- Yeast nutrient, commercial**
Pilot plant scale evaluation for high test molasses fermentation, 72(4), 509.
- Zagrammosoma sp.**
Coffee leaf miner parasitoid in Puerto Rico, 72(2), 255.
- Zea mays**
Estimation of irrigation requirements, 72(2), 277.
N applications and population densities, 72(1), 127.

Zea mays, cv Suresweet

Tres niveles de riego y crecimiento y rendimiento, 72(4), 565.

Zymomonas

Fermentation patterns in high test molasses and blackstrap molasses, 72(3), 475.

Zymomonas mobilis

Batch and fed-batch fermentation of high test molasses, 72(3), 485.

Ethanol production from blackstrap molasses, 72(3), 483.