

SUBJECTS

- Beans, white, green or dry
Mineral losses in hot water and micro-wave blanching, 77(3-4), 129.
- Bemisia tabaci*
Desaerrollo de 4 genotipos de *Phaseolus vulgaris* con diferentes grados de pubescencia, 77(1-2), 61.
- Chicken products
Canned, 77(3-4), 137.
Frozen, 77(3-4), 145.
- Coffee
Fluazifop and quizalofop, grass herbicides in, 77(3-4), 201.
- Cosmopolites sordidus*
Mortality of larvae caused by entomogenous nematodes *Heterorhabditis* spp. 77(1-2), 53.
- Cows, diary
Chopped and ground hay vs. grazing plus concentrates, 77(3-4), 193.
- Cylas formicarius*
Mortality of larvae caused by entomogenous nematode *Heterorhabditis* spp, 77(1-2), 53.
- Dioscorea alata*
Three planting dates and two planting systems, 77(3-4), 161.
Tillage, fertilizer rates and yields, 77(3-4), 153.
- Emesinae
New fossil reduviid from Dominican amber, 77(1-2), 95.
- Euscepes postfasciatus*
Mortality of larvae caused by entomogenous nematodes *Heterorhabditis* spp., 77(1-2), 53.
- Galleria mellonella*
Mortality of larvae caused by entomogenous nematode *Heterorhabditis* spp., 77(1-2), 53.
- Glycine max*
Visual indicator for harvesting immature viable seed, 77(1-2), 33.
- Hay, grass, chopped and ground
Mixed rations vs. grazing plus concentrates, 77(2-4), 193.
- Heteroptera
New fossil reduviid from Dominican amber, 77(1-2), 95.
- Heterorhabditis* spp, entomogenous nematode
Mortality of larvae of *Galleria mellonella*, *Cylas formicarius*, *Euscepes postfasciatus* and *Cosmopolites sordidus*, 77(1-2), 54.
- Leaching, magnesium fertilizers
Puerto Rican Ultisols, 77(3-4), 167.
- Lycopersicon lycopersicum*
Shrimp shell for controlling root node nematode, 77(3-4), 229.
- Meloidogyne incognita*
Exoesqueleto de camarón para controlarlo en tomate, 77(3-4), 229.
- Mg fertilizers
Leaching in Puerto Rican Ultisols, 77(3-4), 167.
- Nematodo nodulador
Exoesqueleto de camarón para controlarlo en tomate, 77(3-4), 229.
- Paleoptoiariola venosa*
New fossil Emesinae, 77(1-2), 95.
- Phaseolus vulgaris*
Desarrollo de *Bemisia tabaci* en 4 genotipos con diferentes grados de pubescencia, 77(1-2), 61.
- Pineapple fields
Grass herbicides, 77(1-2), 45.
- Pineapple pulp, freeze-dried
Comparison of moisture sorption isotherm models, 77(3-4), 113.
- Pseudomonas*
Diseases caused in cultivars in Puerto Rico, 77(3-4), 207.
- Reduviidae
New fossil from Dominican amber, 77(1-2), 95.

- Rums, concentrates
Identification of compounds by gas chromatography and mass spectrometry, 77(1-2), 69.
- Soils, clay
Structural stability, pore size distribution and surface charge properties, 77(1-2), 11.
- Soybean, indeterminate genotypes
Visual indicator for harvesting immature viable seed, 77(1-2), 33.
- Tanier, 3 cultivars
Nutrient uptake, growth and yield, 77(1-2), 1.
- Tomate
Exoesqueleto de camarón para controlar el nematodo nodulador, 77(3-4), 229.
- Tomato production
Chicken manure and chemical properties of a Mollisol, 77(3-4), 181.
- Xanthosoma* spp.
Nutrient uptake, growth and yield, 77(1-2), 1.
- Yams
Three planting dates and two planting systems, 77(3-4), 161.
Tillage, fertilizer rates and yields, 77(3-4), 153.